# $DiaPaste^{\text{TM}} \text{ Root Canal Filling Material}$

# Instructions for Use

Diapaste is a calcium hydroxide paste with barium sulfate, used as a temporary root canal filling material

## Purpose of Use

Aqueous ointment material that temporarily fills the root canal for the following indications:

- Apexification
- Treatment for infected root canal
- Temporary root filling
- Root canal filling for primary teeth
- Disinfection after pulp extirpation treatment
- Vital pulpotomy
- Temporary pulp capping

### **Performance**

- Antibacterial
- Excellent radiopacity
- · Filling microcavities with high flowability
- · Low toxicity and less stimulation to apical tissue
- Stable as it does not solidify nor separate
- Excellent accessibility to the apex
- · High biocompatibility to the periapical tissue

# **Recommended Procedure**

- Preparation before use
- ① Check the product for leaks, discoloration, abnormal condition of the packaging, and expiration date.
- ② A rubber dam is attached to the patient's mouth to protect the mouth. Prepare the necessary accessories for the procedure.
- 3 Be familiar with the warning, side effects, contraindications and instructions.
- How to use
- ① Open the protective cap and attach the disposable tip onto the syringe.
- ② Insert the tip from the root opening to 1/5 of the length of the root canal.

- 3 Push the plunger slowly and gently to inject the contents into the canal.
- ④ Remove excess material in the root canal and proceed with routine restoration procedure.
- ⑤ After use, remove the disposable tip attached to the syringe and attach a new disposable tip.
- 6 Cover the new disposable tip with the provided silicone cap and make sure it is tightly sealed.
- Storage
- ① Keep the silicone cap tightly sealed on the new disposable tip for storage.
- 2 Avoid high temperature and high humidity.
- 3 Do not store near fire.
- 4 Keep out of reach of children.
- $\odot$  Avoid exposure to direct sunlight and store in a dry, cool place. Storage temperature: 2  $^{\circ}$ C  $^{\sim}$  27  $^{\circ}$ C

# Precautions when using

- General Cautions
  - ① This product should only be used by a dentist or a dental professional.
  - ② Do not use on patients with inflammation or allergy to any of this product's components.
  - 3 Avoid contact with skin and eyes. In case of contact, rinse immediately with running water.
  - 4 This product cannot be prescribed for internal use.
  - ⑤ This filling should only remain in the body for 29 days or LESS
- Application Note
  - Do not extrude the product passed the apex.
  - Do not use products that have expired.
  - Do not mix with other products.

# **Expiration**

3 years from date of manufacture

DiaDent Group International #11 – 3871 North Fraser Way Burnaby, BC Canada V5J 5G6

Tel: 1-877-342—3368